



**I AM
TAKING
ON MS**

**I AM PLANNING ON
STARTING A FAMILY**

Rebif® may be an option for treating your relapsing multiple sclerosis (RMS) while planning for a family.

Talk to your healthcare provider to see if Rebif is the right choice for you.

INDICATION

Rebif (interferon beta-1a) is a prescription medicine used to treat relapsing forms of multiple sclerosis, to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults. It is not known if Rebif is safe and effective in children.

IMPORTANT SAFETY INFORMATION

Do not take Rebif if you are allergic to interferon beta, human albumin, or any of the ingredients in Rebif.

Please see additional [**Important Safety Information**](#) throughout and on page 8. See [**Rebif Prescribing Information**](#) and [**Medication Guide**](#) and discuss with your healthcare provider.

**Rebif®**
(interferon beta-1a)
subcutaneous injection



THINKING OF STARTING A FAMILY? THERE'S A LOT TO CONSIDER

If you have relapsing multiple sclerosis (RMS) and want to start a family, it's never been more important to do your homework and plan ahead.

Talk to your healthcare provider early

Your healthcare provider can help you understand how RMS may affect you before, during, and after pregnancy. They'll also guide you on how to manage your RMS treatment during this exciting time.

Ask if Rebif® may be a good choice for you

As you and your healthcare provider review options for treating your RMS while you're planning to have a family, consider Rebif.

The following pages include efficacy data for Rebif, as well as data from a study that looked at pregnancy outcomes for women with RMS who had and had not been exposed to interferon beta in early pregnancy.

This information, plus the Doctor Discussion Guide on page 7 of this brochure may help support a productive discussion with your healthcare provider.

IMPORTANT SAFETY INFORMATION (Cont'd)

Rebif can cause serious side effects. Tell your healthcare provider right away if you have any of the symptoms listed below while taking Rebif.

- **Behavioral health problems including depression and suicidal thoughts.** You may have mood problems including depression (feeling hopeless or feeling bad about yourself), and thoughts of hurting yourself or suicide.

Please see additional [Important Safety Information](#) throughout and on page 8. See [Rebif Prescribing Information](#) and [Medication Guide](#) and discuss with your healthcare provider.

Rebif®
(interferon beta-1a)
subcutaneous injection

Rebif® was proven effective across 3 important treatment goals

The 2-year PRISMS study looked at 560 people with relapsing multiple sclerosis (RMS) to see how they responded to treatment with Rebif. Pregnant women were not allowed to participate in the study, and women who became pregnant during the study were discontinued.

Treatment was given under the skin 3 times a week: 189 people took Rebif 22 mcg, **184 took Rebif 44 mcg**, and **187 took a placebo**. The overall results for people taking Rebif 44 mcg showed:



32% fewer relapses
in 2 years ($P < 0.0001$)

1.73 relapses, on average,
with Rebif (n=184)

2.56 relapses, on average,
with placebo (n=187)



78% fewer new or enlarging
T2 brain lesions
at 2 years* ($P < 0.0001$)

0.5 lesions with Rebif (n=171) | 2.25 lesions with placebo (n=172)



Nearly
2x longer to disability
progression†

21.3 months for Rebif (n=184) | 11.9 months for placebo (n=187)

IMPORTANT SAFETY INFORMATION (Cont'd)

- **Liver problems or worsening of liver problems including liver failure.** Symptoms may include nausea, loss of appetite, tiredness, dark colored urine and pale stools, yellowing of your skin or the white part of your eye, bleeding more easily than normal, confusion, and sleepiness. During your treatment with Rebif you will need to see your healthcare provider regularly and have regular blood tests to check for side effects.

Rebif can cause serious side effects:

- Behavioral health problems including depression and suicidal thoughts
- Liver problems or worsening of liver problems including liver failure
- Serious allergic and skin reactions
- Injection site problems
- Blood problems
- Pulmonary arterial hypertension
- Seizures

The most common side effects of Rebif include:

- Flu-like symptoms
- Stomach pain
- Change in liver blood tests




Although your doctor may determine that you can continue on Rebif until you become pregnant, **you should call your healthcare provider immediately and stop taking Rebif if you become pregnant.**

*Median number of lesions, per patient per scan, based on comparisons from rank-based analysis of variance (ANOVA). Lesions detected with both T1-Gd+ and T2-weighted MRI. Median = a value in an ordered set that has an equal number of values higher and lower.

†Disability progression was defined as an increase of at least 1 point in the Expanded Disability Status Scale (EDSS) that was sustained for at least 3 months.

Please see additional [Important Safety Information](#) throughout and on page 8. See [Rebif Prescribing Information](#) and [Medication Guide](#) and discuss with your healthcare provider.

Rebif®
(interferon beta-1a)
subcutaneous injection



Rebif® has been studied and used widely

With more than
20 years

of combined clinical trial data and
real-world patient experience, Rebif
has a well-established safety profile

150,000

US patients prescribed since approval

IMPORTANT SAFETY INFORMATION (Cont'd)

- **Serious allergic and skin reactions.** Symptoms may include itching, swelling of your face, eyes, lips, tongue or throat, trouble breathing, anxiousness, feeling faint, skin rash, hives, sores in your mouth, or skin blisters and peels.
- **Injection site problems.** Rebif may cause redness, pain, itching or swelling at the place where your injection was given. Call your healthcare provider right away if an injection site becomes swollen and painful or the area looks infected. You may have a skin infection or an area of severe skin damage (necrosis) requiring treatment by a healthcare provider.

Common side effects

Common adverse events have been consistent across PRISMS Study. It's important to inform your healthcare provider of any side effects that you experience. There may be things you can do to help manage them.

Common side effects seen in the PRISMS study

This chart tells you the most common side effects and the percentage of people who experienced them during the study.

Side effects	Rebif 44 mcg (%)	Placebo (%)
Headache	70%	63%
Flu-like symptoms	59%	51%
Fatigue	41%	36%
Fever	28%	16%
Injection-site reactions	92%	39%

These are not all the possible side effects associated with Rebif.

PRISMS = **P**revention of **R**elapses and **D**isability by **I**nterferon β -1a **S**ubcutaneously in **M**ultiple **S**clerosis.

Please see additional [Important Safety Information](#) throughout and on page 8. See [Rebif Prescribing Information](#) and [Medication Guide](#) and discuss with your healthcare provider.





The pregnancy registry

In a study conducted in Finland from 1996-2014 and Sweden from 2005-2014, data were collected about 2,831 pregnancy outcomes from women with MS.

In early pregnancy, some women were exposed to interferon beta (n=797 pregnancies) while others were not exposed to any disease-modifying drugs (n=1,647 pregnancies).

Understanding the Registry Data

Of the 797 pregnancies that were only exposed to interferon beta, including Rebif® (interferon beta-1a), the data:

DID NOT IDENTIFY

an increased drug-associated risk of major birth defects among the women exposed to interferon beta in early pregnancy.

DID NOT IDENTIFY

an increased risk of miscarriage or ectopic pregnancy. However, limitations in obtaining complete data capture for these outcomes made the interpretation of these findings more difficult.

SELECTED IMPORTANT SAFETY INFORMATION


Before you take Rebif, tell your healthcare provider if you have or have had any of the following conditions:

- are pregnant or plan to become pregnant. It is not known if Rebif will harm your unborn baby. Tell your healthcare provider if you become pregnant during your treatment with Rebif
- are breastfeeding or plan to breastfeed. Rebif may pass into your breastmilk. Talk with your healthcare provider about the best way to feed your baby if you take Rebif.

Understanding the registry data

Pregnancy outcomes (live births without birth defects) of women in the Nordic MS Pregnancy Registry Study and the US general population:


Proportion of live births without birth defects



Nordic MS group
not exposed to
interferon beta
(n=1,506)*



Nordic MS group
exposed only to
interferon beta
(n=683)*



US general
population

*The data shown above are from an analysis that differs from the analysis reported in the Prescribing Information for Rebif.



Although your doctor may determine that you can continue on Rebif until you become pregnant, **you should call your healthcare provider immediately and stop taking Rebif if you become pregnant.**

Please see additional [Important Safety Information](#) throughout and on page 8. See [Rebif Prescribing Information](#) and [Medication Guide](#) and discuss with your healthcare provider.

Rebif®
(interferon beta-1a)
subcutaneous injection

RMS: before, during, and after pregnancy

If you are planning to get pregnant, it is important to talk with your healthcare provider about managing your risk of relapsing multiple sclerosis (RMS) relapses (flare-ups) before, during, and after pregnancy.



Before pregnancy

Rebif® may be an appropriate choice to treat your RMS while you plan for pregnancy. If you're thinking about becoming pregnant, ask your healthcare provider whether Rebif may be an option for you.



During pregnancy

Tell your healthcare provider right away if you become pregnant. Although the study data did not identify an increased drug-associated risk of major birth defects with interferon beta use during early pregnancy, it is not known if Rebif may harm your unborn baby.




After pregnancy

Your healthcare provider can help you decide when it's the right time to start (or restart) Rebif after giving birth. Talk to them about the risk of relapses after birth and how soon you should consider starting treatment.

IMPORTANT SAFETY INFORMATION (Cont'd)

- **Blood problems.** Rebif can affect your bone marrow and cause low red and white blood cell and platelet counts. In some people, these blood cell counts may fall to dangerously low levels. If your blood cell counts become very low, you can get infections and problems with bleeding and bruising. Your healthcare provider may ask you to have regular blood tests to check for blood problems.



Hoping to breastfeed your baby?

You may also wonder about breastfeeding your baby postpartum. There is limited published data on this topic. Some data has shown interferon may be present at low levels in breastmilk. Other data has shown that it is poorly absorbed into the baby's bloodstream. Talk with your healthcare provider about the best way to feed your baby if you take Rebif.



Could Rebif be right for you?

Use the discussion guide on the following page of this brochure to help start a conversation with your healthcare provider.

Please see additional [Important Safety Information](#) throughout and on page 8. See [Rebif Prescribing Information](#) and [Medication Guide](#) and discuss with your healthcare provider.

Rebif®
(interferon beta-1a)
subcutaneous injection



Talking with your doctor

Consider these questions to help guide discussions with your healthcare provider about pregnancy and relapsing multiple sclerosis (RMS).

☐ Can I take Rebif® when trying to get pregnant?

☐ Are pregnancy symptoms different for women with RMS?

☐ What can I expect to happen with relapses when I am pregnant?

☐ How can I protect myself against relapses after having my baby?

☐ What should I do about an RMS treatment plan after my baby comes?

INDICATION

Rebif® (interferon beta-1a) is a prescription medicine used to treat relapsing forms of multiple sclerosis, to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults. It is not known if Rebif is safe and effective in children.

☐ What are my options for feeding my baby?

☐ Where can I get more information about pregnancy and RMS?

Notes/your own questions

Please see additional [Important Safety Information](#) throughout and on page 8. See [Rebif Prescribing Information](#) and [Medication Guide](#) and discuss with your healthcare provider.

US-REB-00451 10/23

 **Rebif®**
(interferon beta-1a)
subcutaneous injection

INDICATION AND IMPORTANT SAFETY INFORMATION for REBIF® (interferon beta-1a) subcutaneous injection

INDICATION

Rebif® (interferon beta-1a) is a prescription medicine used to treat relapsing forms of multiple sclerosis, to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults. It is not known if Rebif is safe and effective in children.

IMPORTANT SAFETY INFORMATION

Do not take Rebif if you are allergic to interferon beta, human albumin, or any of the ingredients in Rebif.

Rebif can cause serious side effects. Tell your healthcare provider right away if you have any of the symptoms listed below while taking Rebif.

- **Behavioral health problems including depression and suicidal thoughts.** You may have mood problems including depression (feeling hopeless or feeling bad about yourself), and thoughts of hurting yourself or suicide.
- **Liver problems or worsening of liver problems including liver failure.** Symptoms may include nausea, loss of appetite, tiredness, dark colored urine and pale stools, yellowing of your skin or the white part of your eye, bleeding more easily than normal, confusion, and sleepiness. During your treatment with Rebif you will need to see your healthcare provider regularly and have regular blood tests to check for side effects.
- **Serious allergic and skin reactions.** Symptoms may include itching, swelling of your face, eyes, lips, tongue or throat, trouble breathing, anxiousness, feeling faint, skin rash, hives, sores in your mouth, or skin blisters and peels.
- **Injection site problems.** Rebif may cause redness, pain, itching or swelling at the place where your injection was given. Call your healthcare provider right away if an injection site becomes swollen and painful or the area looks infected. You may have a skin infection or an area of severe skin damage (necrosis) requiring treatment by a healthcare provider.
- **Blood problems.** Rebif can affect your bone marrow and cause low red and white blood cell and platelet counts. In some people, these blood cell counts may fall to dangerously low levels. If your blood cell counts become very low, you can get infections and problems with bleeding and bruising. Your healthcare provider may ask you to have regular blood tests to check for blood problems.
- **Pulmonary arterial hypertension.** Pulmonary arterial hypertension can occur with interferon beta products, including Rebif. Symptoms may include new or increasing fatigue or shortness of breath. Contact your healthcare provider right away if you develop these symptoms.
- **Seizures.** Some people have had seizures while taking Rebif.

Before you take Rebif, tell your healthcare provider if you have or have had any of the following conditions:

- mental illness, including depression and suicidal behavior
- liver problems
- bleeding problems or blood clots
- low blood cell counts
- seizures (epilepsy)
- thyroid problems
- drink alcohol
- are pregnant or plan to become pregnant. It is not known if Rebif will harm your unborn baby. Tell your healthcare provider if you become pregnant during your treatment with Rebif
- are breastfeeding or plan to breastfeed. Rebif may pass into your breastmilk. Talk with your healthcare provider about the best way to feed your baby if you take Rebif.

Tell your healthcare provider about all medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

The most common side effects of Rebif include:

- flu-like symptoms. You may have flu-like symptoms when you first start taking Rebif. You may be able to manage these flu-like symptoms by taking over-the-counter pain and fever reducers. For many people, these symptoms lessen or go away over time. Symptoms may include muscle aches, fever, tiredness, and chills
- stomach pain
- change in liver blood tests

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of Rebif. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or visit www.fda.gov/medwatch.

See [Prescribing Information](#) and [Medication Guide](#).

 **Rebif®**
(interferon beta-1a)
subcutaneous injection



 **Rebif**[®]
*(interferon beta-1a)
subcutaneous injection*

Ask your healthcare provider if Rebif[®] may be an option for managing your RMS while planning for a family.

For more information, visit rebif.com

Please see additional **Important Safety Information** throughout and on page 8. See **Rebif Prescribing Information** and **Medication Guide** and discuss with your healthcare provider.

**EMD
SERONO**

The people depicted in this brochure are not real patients and are not sponsored by EMD Serono, Inc. Rebif[®] is a registered trademark of Merck KGaA, Darmstadt, Germany, or its affiliates. EMD Serono, Inc. is the healthcare business of Merck KGaA, Darmstadt, Germany in the U.S. and Canada. ©2023 Merck KGaA, Darmstadt, Germany or its affiliates. All rights reserved. US-REB-00451 10/23